

AD \_\_\_\_\_

Award Number: W81XWH-06-1-0172

TITLE: On the Pathophysiology of Cutaneous Café-au-lait Lesions in Neurofibromatosis and the Role of Keratinocyte and/or Fibroblast-Synthesized Cytokines

PRINCIPAL INVESTIGATOR: Raymond E. Boissy, Ph.D.

CONTRACTING ORGANIZATION: University of Cincinnati  
Cincinnati, OH 45267

REPORT DATE: January 2007

TYPE OF REPORT: Final

PREPARED FOR: U.S. Army Medical Research and Materiel Command  
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;  
Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. <b>PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.</b>					
1. REPORT DATE (DD-MM-YYYY) January 2007		2. REPORT TYPE Final		3. DATES COVERED (From - To) 15 Dec 2005 – 14 Dec 2006	
4. TITLE AND SUBTITLE  On the Pathophysiology of Cutaneous Café-au-lait Lesions in Neurofibromatosis and the Role of Keratinocyte and/or Fibroblast-Synthesized Cytokines				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER W81XWH-06-1-0172	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Raymond E. Boissy, Ph.D.  E-Mail: <a href="mailto:dqku{tgB weo ckr0eQf w">dqku{tgB weo ckr0eQf w</a>				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)  University of Cincinnati Cincinnati, OH 45267				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT:  Fibroblasts derived from the CALM lesion of NF1 are capable of inducing melanin synthesis by normal human melanocytes as demonstrated in both an in vitro and in vivo model system suggesting that undefined cytokines are synthesized by NF1 fibroblasts putatively resulting in CALM.					
15. SUBJECT TERMS Skin, café-au-lait, pigmentation, melanocytes					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON
a. REPORT	b. ABSTRACT	c. THIS PAGE			USAMRMC
U	U	U	UU	6	19b. TELEPHONE NUMBER (include area code)

	<b>Page</b>
<b>Introduction.....</b>	<b>4</b>
<b>Body.....</b>	<b>4</b>
<b>Key Research Accomplishments.....</b>	<b>5</b>
<b>Conclusion.....</b>	<b>6</b>

## **INTRODUCTION**

Mutations at the *NF1* locus predisposes neurofibromatosis type 1 (NF1) patients to skin pigmentary abnormalities: hyperpigmented café-au-lait macules (CALM) and axillary & inguinal freckling. The pathophysiology of these *NF1*-related abnormalities has yet to be elucidated. Much controversial data exists pertaining to the etiology of these pigmentary lesions, however the consensus suggests that the melanocyte population in CALM is hyperactive resulting in increased pigment synthesis. However, melanocytes cultured in isolation from skin biopsies of NF1 patients demonstrated no, or at most minimal, increase in melanization. This suggests that extracellular factors (i.e., cytokines) in the NF1 skin may be the important elements inducing hyperactivity of the melanocytes resulting in CALM. We proposed that an NF1 mutation, and/or loss of heterozygosity, causes aberrant production of secretory cytokines by the NF1 fibroblast of the skin that ultimately influences the melanocytes resulting in CALM. This study was designed to understand the etiology of CALM in NF1 by assessing interactions between melanocytes (M) and fibroblasts (Fb) in culture derived from NF1 patients.


## **BODY**

To test the hypothesis proposed above, we initially assessed the proliferation and melanization of cultured M derived from CALM-lesional sites and normal skin under normal and promotional (i.e., increased growth factors) conditions. Subsequently we assessed serum-free, conditioned media from cultured Fb derived from CALM-lesional sites in comparison to Fb derived from normal individuals, for the ability to stimulate proliferation and melanogenesis on cultured normal M as well as melanocytes derived from CALM-lesional sites in several patients with NF1.

## KEY RESEARCH ACCOMPLISHMENTS


### Proliferation and response to growth factors were similar between NHM and NF1

melanocytes. – Melanocyte cultures were developed from 5 patients with NF1 and 5 unaffected individuals. Proliferation of cultures were assess in media with 1, 2 or 5 X growth factors (GF) added. No statistical differences were observed.



### Induction of melanization by growth factors was enhanced in NF1 melanocytes compared

to NHM. – Cultures described immediately above were assessed for melanin content. NF1 melanocytes exhibited a statistically significant increase in melanin by GF.




### Proliferation in response to conditioned media from NF1-fibroblasts was similar between

NHM and NF1 melanocytes. – Fibroblasts cultures were developed from 5 patients with NF1 and 5 unaffected individuals. Conditioned media from these cultures was then combined with normal melanocyte growth media at various ratios for 10 days and proliferation subsequently assessed. No significant differences were observed among the various groups (data not shown).

### Induction of melanization by conditioned media from NF1-fibroblasts was enhanced in

NF1 melanocytes compared to NHM. Cultures described immediately above were assessed for melanin content. NHM exhibited a statistically significant increase in melanin induced by NF1 derived Fb as oppose to normal Fb.



## **CONCLUSION**

Fibroblasts derived from the CALM lesion of NF1 are capable of inducing melanin synthesis by normal human melanocytes as demonstrated in an *in vitro* model system suggesting that undefined cytokines are synthesized by NF1 fibroblasts putatively resulting in CALM.